

*This copy is for your personal, non-commercial use only.*

**If you wish to distribute this article to others**, you can order high-quality copies for your colleagues, clients, or customers by [clicking here](#).

**Permission to republish or repurpose articles or portions of articles** can be obtained by following the guidelines [here](#).

**The following resources related to this article are available online at  
www.sciencemag.org (this information is current as of February 27, 2013):**

**Updated information and services**, including high-resolution figures, can be found in the online version of this article at:

<http://www.sciencemag.org/content/339/6122/913.full.html>

This article **cites 13 articles**, 2 of which can be accessed free:

<http://www.sciencemag.org/content/339/6122/913.full.html#ref-list-1>

This article appears in the following **subject collections**:

Biochemistry

<http://www.sciencemag.org/cgi/collection/biochem>

1-independent pyroptosis, as well as caspase-1-dependent cytokine production, in response to Gram-negative bacteria (6–10).

Aachoui *et al.* establish that caspase-11 has a protective role during infection with cytosolic bacteria. The authors show that mice lacking both caspase-1 and -11 (*Casp1<sup>-/-</sup>**Casp11<sup>-/-</sup>*) succumb to infection by the cytosol-adapted Gram-negative pathogen *Burkholderia thailandensis*. By contrast, mice lacking functional inflammasomes (through the deletion of genes encoding the inflammasome components NLRP3, NLRC4, and ASC), and that also have impaired IL-1 $\beta$  secretion, were resistant to *B. thailandensis*, indicating that resistance does not depend on cytokine secretion. The *Casp1<sup>-/-</sup>**Casp11<sup>-/-</sup>* macrophages did not activate cell death by pyroptosis in response to the bacterial infection, unlike macrophages lacking inflammasomes. Aachoui *et al.* demonstrate that caspase-11 is primarily responsible for pyroptosis and that *Casp1<sup>-/-</sup>* mice recapitulate the lethal phenotype of *Casp1<sup>-/-</sup>**Casp11<sup>-/-</sup>* mice with *B. thailandensis* infection. Caspase-11-dependent pyroptosis resembled caspase-1-dependent pyroptosis, as judged by membrane damage as well as glycine sensitivity (glycine blocks ion fluxes in damaged eukaryotic cells, thereby preventing swelling and lysis in pyroptosis triggered by caspase-1).

Aachoui *et al.* further support their finding that caspase-11 protects against cytosolic bacteria by using the vacuole-adapted pathogen, *Salmonella typhimurium*. A mutant strain,  $\Delta$ sifA, localizes to the cytosol because it does not secrete SifA, a “type 3 secreted effector” protein that maintains the integrity of *Salmonella*-containing vacuole (11). Only in the presence of caspase-11 can the wild-type bacteria substantially outcompete the  $\Delta$ sifA mutant. This implies that wild-type *S. typhimurium* residing in the vacuole that are not exposed to caspase-11 are protected, whereas the cytosol-localized  $\Delta$ sifA mutant is vulnerable to caspase-11 detection and subsequent elimination by pyroptosis.

It is noteworthy that the authors observe induction of caspase-11 expression after priming by bacterial lipopolysaccharide (LPS) or interferon- $\gamma$  (IFN- $\gamma$ ) a cytokine that is critical for immunity against intracellular bacterial infections. Caspase-11 may be constitutively active upon its expression (10). However, the finding that bacterial entry to the cytosol is required for induction of pyroptosis suggests that other signals are required for caspase-11 to execute its function. Whether caspase-11 undergoes post-translational modifications or subcellular

targeting following bacterial escape into the cytosol is not yet known.

Caspase-11 is now recognized as an important part of the inflammatory arsenal that can be either protective (3) or damaging depending on the physiological context of its activity (6, 7). Yet, the study of Aachoui *et al.* raises two major questions. What is the signal(s) that activate caspase-11? A “noncanonical inflammasome” has been proposed to activate caspase-11 (6), but no molecular players have yet been identified. And what are the substrates of caspase-11 that promote pyroptosis? These gaps in understanding of caspase-11 function will be important topics for future studies.

There is a dynamic interplay between caspases and bacterial pathogens. Some bacteria use caspases to promote their pathogenesis. For example, *Listeria monocytogenes* exploits caspase-7 to repair bacteria-induced damage to the plasma membrane to preserve its replicative niche (12). Similarly, *S. typhimurium* exploits caspase-3 to cleave and activate a virulence factor, SipA (13). By contrast, other pathogens inhibit inflammasome activation to attenuate the immune response. Bacterial factors from *Mycobacterium tuberculosis* and *Yersinia pestis* promote their virulence by inhibiting inflam-

masome activation, thereby dampening their host's immune response (14, 15). In the case of *Y. pestis*, its type 3 secreted effector, YopM, sequesters caspase-1 into the nucleus where it can no longer interact with the inflammasome (14). It remains to be elucidated whether some cytosol-adapted pathogens, like *L. monocytogenes* and *Shigella flexneri*, have devised strategies to evade or exploit caspase-11-dependent pyroptosis.

## References

1. M. Cemma, J. H. Brumell, *Curr. Biol.* **22**, R540 (2012).
2. M. J. Davis, B. Gregorka, J. E. Gestwicki, J. A. Swanson, *J. Immunol.* **189**, 4488 (2012).
3. Y. Aachoui *et al.*, *Science* **339**, 975 (2013); 10.1126/science.1230751.
4. A. S. Yazdi, G. Guarda, M. C. D'Ombrain, S. K. Drexler, *J. Innate Immun.* **2**, 228 (2010).
5. E. A. Miao, J. V. Rajan, A. Aderem, *Immunol. Rev.* **243**, 206 (2011).
6. N. Kayagaki *et al.*, *Nature* **479**, 117 (2011). C.
7. P. Broz *et al.*, *Nature* **490**, 288 (2012).
8. C. L. Case *et al.*, *Proc. Natl. Acad. Sci. U.S.A.* **110**, 1851 (2013).
9. P. Gurung *et al.*, *J. Biol. Chem.* **287**, 34474 (2012).
10. V. A. Rathinam *et al.*, *Cell* **150**, 606 (2012).
11. C. R. Beuzón *et al.*, *EMBO J.* **19**, 3235 (2000).
12. S. K. Cassidy *et al.*, *PLoS Pathog.* **8**, e1002628 (2012).
13. C. V. Srikanth *et al.*, *Science* **330**, 390 (2010).
14. C. N. LaRock, B. T. Cookson, *Cell Host Microbe* **12**, 799 (2012).
15. S. S. Master *et al.*, *Cell Host Microbe* **3**, 224 (2008).

10.1126/science.1235639

## BIOCHEMISTRY

# Integrative Structural Biology

Andrew B. Ward,<sup>1</sup> Andrej Sali,<sup>2</sup> Ian A. Wilson<sup>1,3</sup>

Integrative approaches using data from a wide variety of methods are yielding model structures of complex biological assemblies.


**B**iological assemblies and machines often elude structural characterization, hampering our understanding of how they function, how they evolved, and how they can be modulated. A number of macromolecular assemblies have been reconstructed over the years by piecemeal efforts, such as fitting high-resolution crystal structures of individual components into lower-resolution

electron microscopy (EM) reconstructions of the entire complex (1). Although notable successes have been achieved in this way, ambiguous or conflicting models can still arise (2–4). Thus, structural and computational biologists have been looking for new ways to put all of the pieces back together. Sophisticated integrative approaches are being developed (5, 6) that combine information from different types of experiments, physical theories, and statistical analyses to compute structural models of multicomponent assemblies and complex biological systems.

In addition to the conventional biophysical techniques of x-ray crystallography, nuclear magnetic resonance (NMR) spectroscopy, EM, and small-angle x-ray scattering (SAXS), a growing number of experimental methods can also provide valuable infor-

<sup>1</sup>Department of Integrative Structural and Computational Biology, International AIDS Vaccine Initiative (IAVI) Neutralizing Antibody Center, and Center for HIV/AIDS Vaccine Immunology and Immunogen Discovery (CHAVI-ID), Scripps Research Institute, La Jolla, CA 92037, USA.

<sup>2</sup>Department of Bioengineering and Therapeutic Sciences, Department of Pharmaceutical Chemistry, and California Institute for Quantitative Biosciences (QB3), University of California, San Francisco, CA 94158, USA. <sup>3</sup>Skaggs Institute for Chemical Biology, Scripps Research Institute, La Jolla, CA 92037, USA. E-mail: abward@scripps.edu



**Complex structure solutions.** Models of macromolecules and their complexes can be constructed by combining different types of information generated by various experimental and theoretical techniques (gray box). The data are converted into spatial restraints, which are combined into a scoring function that guides sampling algorithms to obtain a detailed structural model.

mation about the structures and dynamics of proteins and their assemblies. These methods include sequence comparisons of related proteins, copurification, hydrogen-deuterium exchange mass spectrometry (HDXMS), single-molecule fluorescence, atomic force microscopy, analytical spectroscopy (both electron paramagnetic resonance and double electron-electron resonance), light scattering, chemical cross-linking, and mutagenesis (see the figure).

The individual pieces of data gathered using different techniques can provide invaluable restraints on the conformation, position, and orientation of the components in an assembly or biological system (5). Relative to the use of any single set or type of data, simultaneous use of all such restraints can markedly improve the accuracy, precision, and completeness of a model, especially when high-resolution structural data on the entire complex are not available.

Because of the many degrees of freedom in macromolecular structures and the difficulty of combining disparate data, models must be computed with algorithms that sam-

ple as many potential solutions as possible given the computing power available. These algorithms are driven by a scoring function consisting of the individual spatial restraints and are analogous to methods used in x-ray crystallography and NMR spectroscopy, which also generate models by minimizing differences between experimental data and data calculated from a model. Assessing how to best combine and weigh different types of data from multiple sources is a prerequisite for constructing structural models of increasingly larger and more dynamic macromolecular complexes.

A useful test of a model is whether it explains all data points within their own error bars and whether the entire data set is redundant, meaning that a subset of the data can be omitted without any significant impact on the model. In such a case, the confidence in the model, the data, and the parameters used for modeling can be high. When a subset of the data points cannot be satisfied by a single model because the data were collected from a heterogeneous sample and/or the data are noisy, more sophisticated methods for com-

bining individual restraints are needed. In such cases, emphasis is placed on evaluating models in an objective manner, using Bayesian (7) and other statistical methods that explicitly take into account the noise in the data and/or multiple structural states in the sample.

Integrative, restraint-based approaches can be used whenever a challenging structural biology problem is encountered, from an individual protein to a small macromolecular machine to a large multicomponent cellular assembly. Thus, integrative approaches span wide resolution ranges and bridge observations made from the atomic to the cellular level. The following three examples illustrate the power of these new methods in generating models at different levels of resolution.

Some of the most successful applications of integrative approaches have resulted from combining sparse experimental observations with computation to generate atomic-level models of macromolecules. Rosetta (8), a platform for modeling protein structures, works by exhaustive calculations under a set of assumptions about the underlying geometry and chemistry of peptides. These assumptions reduce the nearly infinite sampling necessary to fold a one-dimensional sequence of amino acids into a three-dimensional shape. Experimental restraints from NMR (9) or EM (10) can further narrow the search and help to converge on more accurate models. For example, Loquet *et al.* used solid-state NMR, EM, and Rosetta to build an atomic-level model of the bacterial type III secretion needle used to inject its proteins into host cells (11). The model revealed details of the supramolecular interfaces of the component protomers, providing a structural understanding of this machine that had eluded characterization by single techniques.

Two recent independent studies of the molecular architecture of the 26S proteasome exemplify the value of integrative approaches for medium-resolution structures. Lander *et al.* combined EM reconstructions and x-ray crystal structures (12), whereas Lasker *et al.* used restraints from a variety of data sets (EM, x-ray crystallography, chemical cross-linking, and proteomics) and employed the Integrative Modeling Platform package (6, 13) to build an almost identical model of the 26S proteasome (14). Lasker *et al.*'s model was further tested by systematically removing some input data, recalculating a model, and assessing it against the omitted data. Although neither model resolved all interactions at an atomic level, they provided a detailed understanding of the arrangement of the component subunits and were therefore

extremely informative about the evolution and function of the 26S proteasome.

At low resolution, chromatin has also been modeled through integrative approaches. In this way, Duan *et al.* constructed a three-dimensional model of the yeast genome (15), uncovering the topology and spatial relationships of different chromosomal elements. For this study, the restraints were garnered from cross-linking, restriction enzyme digestion, ligation, and deep sequencing, thereby revealing the three-dimensional structure of the genome at a level of detail not accessible to any conventional imaging method typically used to study assemblies of this size. These inferential, cellular-scale approaches enable comparison of normal and aberrant cells and may eventually serve an important diagnostic role in medicine.

As integrative methods evolve, structural models can then be revisited and new data incorporated, so that the model can be contin-

uously improved and revised using the latest information (13). Integrative software tools should therefore be flexible enough to incorporate new data and/or restraints. The Protein Data Bank ([www.pdb.org](http://www.pdb.org)) is facilitating this process by acting as a curator for a variety of structural data from different methods as well as models based on these data.

Any experimental observation can in principle be converted into a restraint for building ever more complex models. The reach and impact of structural biology can thus be extended to a wider and more diverse audience. Using these new computational and bioinformatics approaches to collect and integrate diverse pieces of structural and experimental data, Humpty Dumpty can be put back together again.

#### References and Notes

- G. C. Lander, H. R. Saibil, E. Nogales, *Curr. Opin. Struct. Biol.* **22**, 627 (2012).
- H. W. Wang *et al.*, *Nat. Struct. Mol. Biol.* **16**, 1148 (2009).
- P. W. Lau *et al.*, *Nat. Struct. Mol. Biol.* **19**, 436 (2012).
- R. Henderson *et al.*, *Structure* **20**, 205 (2012).
- F. Alber, F. Förster, D. Korkin, M. Topf, A. Sali, *Annu. Rev. Biochem.* **77**, 443 (2008).
- D. Russel *et al.*, *PLoS Biol.* **10**, e1001244 (2012).
- W. Rieping, M. Habeck, M. Nilges, *Science* **309**, 303 (2005).
- [www.rosettacommons.org](http://www.rosettacommons.org)
- N. G. Sgourakis *et al.*, *J. Am. Chem. Soc.* **133**, 6288 (2011).
- F. DiMaio, M. D. Tyka, M. L. Baker, W. Chiu, D. Baker, *J. Mol. Biol.* **392**, 181 (2009).
- A. Loquet *et al.*, *Nature* **486**, 276 (2012).
- G. C. Lander *et al.*, *Nature* **482**, 186 (2012).
- <http://integrativemodeling.org>
- K. Lasker *et al.*, *Proc. Natl. Acad. Sci. U.S.A.* **109**, 1380 (2012).
- Z. Duan *et al.*, *Nature* **465**, 363 (2010).

**Acknowledgments:** Supported by the Protein Structure Initiative of the National Institute of General Medical Sciences, the CHAVI-ID and HIVRAD (HIV Vaccine Research and Design) programs of the National Institute of Allergy and Infectious Diseases, and IAVI.

10.1126/science.1228565

#### PLANT SCIENCE

## Preservation of Recalcitrant Seeds

Christina Walters,<sup>1</sup> Patricia Berjak,<sup>2</sup> Norman Pammerter,<sup>2</sup> Kathryn Kennedy,<sup>3</sup> Peter Raven<sup>4</sup>

**C**oncerns about the rapid erosion of plant diversity have spawned a host of seed-banking initiatives (1). These repositories provide critical germ plasm needed to understand, maintain, and manage natural variation within and among species (2). However, numerous plant species and much of the humid tropics are underserved in these endeavors because of the perceived problem of seed recalcitrance (3). About 75 to 80% of angiosperm species (4, 5) produce orthodox seeds that can survive drying and prolonged storage at -20°C. By contrast, 5 to 10% of angiosperm species produce recalcitrant seeds that do not survive desiccation (3) and are killed in the freezer when ice crystals form. How can their preservation be ensured?

These seeds are called recalcitrant because they die rapidly when treated under the same genebanking conditions used to store orthodox seeds. Recalcitrant seeds,

like most living organisms but unlike orthodox seeds, need water to survive. By nature, recalcitrant seeds are short-lived; they either germinate or are eaten by animals in the wild (3). "Intermediate" seeds, produced by 10 to 15% of angiosperm species (4), withstand sufficient dehydration to prevent formation of lethal ice; nevertheless, their seeds are short-lived in the freezer for unknown reasons (6).

The misperception that recalcitrant seeds cannot be stored arises from the assumption that there is only one storage strategy at our disposal—standard freezers—to manage the wide variation in storage physiology exhibited by plant seeds. In reality, long-term storage of recalcitrant and intermediate seeds (or seed parts) is possible with cryogenic technologies (7–9). Because cryogenic storage requires specialized infrastructure and personnel and is costly, most genebanks, even some of the newer ones, invest only in freezer storage.

Cryopreservation involves storage at ultralow temperature, often in liquid nitrogen (-196°C) (7–9). Rapidly advancing methods can be used to essentially stop water from freezing within recalcitrant seed cells and obviate lethal freezing damage (3, 7–9). The technology requires care-

**Cryogenic technologies help to preserve plant biodiversity in seed banks, particularly in the tropics.**

fully dehydrating tissues and cooling them extremely rapidly (7). Small samples are required to obtain diffusion and heat transfer rates needed to prevent ice formation (7).

The problem is that recalcitrant seeds tend to be large compared with orthodox or intermediate seeds (see the figure) (5). In fact, most recalcitrant seeds are far too large to be rapidly dehydrated or cooled effectively when exposed to liquid nitrogen. The major breakthrough that led to successful cryopreservation of recalcitrant germ plasm was the ability to surgically dissect out the growing portion of the seed (termed the embryonic axis) and germinate it in vitro (10). Recovery may be enhanced by exposing embryonic axes to cryoprotectants (substances that protect against dehydration and freezing damage) (8, 9) and reducing stress-related free radical-mediated damage (11).

Cryogenic technology has been successfully refined for embryonic axes of a spectrum of temperate recalcitrant-seeded species (8). Intermediate seeds are smaller and drier and do not even require these exacting preparations. However, unmet challenges still limit successful cryopreservation of embryonic axes of some recalcitrant seeds from the tropics and subtropics (11). The seeds of these species have embryonic

<sup>1</sup>U.S. Department of Agriculture—Agricultural Research Service National Center for Genetic Resources Preservation, Fort Collins, CO 80521, USA. <sup>2</sup>Plant Germplasm Conservation Research, School of Life Sciences, University of KwaZulu-Natal, Durban, 4001 South Africa. <sup>3</sup>The Center for Plant Conservation, St. Louis, MO 63166-0299, USA. <sup>4</sup>Missouri Botanical Garden, St. Louis, MO 63166-0299, USA. E-mail: christina.walters@ars.usda.gov